DOCUMENT RESUME

ED 243 926 TM 840 231

AUTHOR Federico, Pat-Anthony
TITLE Spatial Ability and Cerebral Sensory Interaction.
PUB DATE Apr 84
NOTE 28p.; Paper presented at the Annual Meeting of the American Educational Research Association (68th, New Orleans, LA, April 23-27, 1984).
PUB TYPE Speeches/Conference Papers (150) -- Reports - Research/Technical (143)

EDRS PRICE MF01/PC02 Plus Postage.
DESCRIPTORS Adults; Cognitive Ability; *Cognitive Processes; Electroencephalography; Enlisted Personnel; *Males; *Neurological Organization; *Sensory Integration; *Spatial Ability; *Visual Stimuli
IDENTIFIERS Brain Hemispheres; *Cerebral Sensory Interaction

ABSTRACT

To provide converging support that the proper integration of analog and propositional representational systems is associated with spatial ability, the visual, auditory, and bimodal brain event-related potentials were recorded from 50 right-handed Caucasian males. Sensory interaction indices were derived for these subjects who had taken the Surface Development Test of spatial ability. Product-moment correlations were computed between sensory interaction indices for eight cerebral sites and spatial ability test scores. Sensory interaction for left and right hemispheric regions was significantly related to spatial ability. As sensory suppression lessened, spatial ability increased. The results substantiated the theory that the visual-imaginal-analog and the auditory-verbal-propositional representational systems are implicated in spatial ability. The extent to which the cortex can inhibit or attenuate the interaction or integration between these dual symbol systems is associated with complicated spatial task performance. (Author)

****
* Reproductions supplied by EDRS are the best that can be made *
* from the original document. *
****

ED243926

Spatial Ability and Cerebral Sensory Interaction

U.S. DEPARTMENT OF EDUCATION
NATIONAL INSTITUTE OF EDUCATION
EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

☒ This document has been reproduced as received from the person or organization originating it.

☐ Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official NIE position or policy.

Pat-Anthony Federico

Navy Personnel Research and Development Center

San Diego, California 92152

"PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY

P-A. Federico

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)."

Paper presented at the meeting of the American Educational Research Association, New Orleans, April, 1984

Opinions and assertions contained herein are those of the author and are not to be construed as official or reflecting views of the Department of the Navy or Defense.

TM 810 231

Abstract

To provide converging support that the proper integration of analog and propositional representational systems is associated with spatial ability, visual, auditory, and bimodal brain event-related potentials were recorded from 50 right-handed Caucasian males. Sensory interaction indices were derived for these subjects who had taken the Surface Development Test of spatial ability. Product-moment correlations were computed between sensory interaction indices for eight cerebral sites and spatial ability test scores. Sensory interaction for left and right hemispheric regions was significantly related to spatial ability. As sensory suppression lessen, spatial ability increased. The results substantiated the theory that the visual-imaginal-analog and the auditory-verbal-propositional representational systems are implicated in spatial ability. The extent to which the cortex can inhibit or attenuate the interaction or integration between these dual symbol systems is associated with complicated spatial task performance.

An important epistemological dispute has centered on the nature of the mental representations intrinsic to spatial ability. The essential issue has focused on whether it is necessary to propose special types of cognitive processes for spatial information in addition to those associated with verbal information. The amount of match or isomorphism between the internal representations and their external referents has been used to differentiate between analog and propositional processes. Analog representational processes preserve the relational structure of external objects, and propositional representational processes do not. An internal transformation is considered analog when the intermediate phases in the process have an isomorphic or one-to-one correspondence with the intermediate phases in the external transformation of the object (Shepard & Cooper, 1982).

There are three attributes that contribute to the definition of propositional representation (Anderson, 1978): (1) it is abstract not dependent on the particular characteristics of its instantiation, (2) it has truth value in that it is either true or false, and (3) it has rules that govern its formation so that decisions can be made regarding how well it is structured. Given these criteria, it seems reasonable to propose that implicit propositional systems underlie natural languages, mathematics, computer programs, and semantic networks. An analog representation does not convey these features. However, it does imply an isomorphism, or one-to-one mapping, between a mental representation and its corresponding real referent, e.g., a road map corresponds to the lay out of streets and highways, a building corresponds to its architectural plans, a football play on the field corresponds to the coach's diagram of it. Analog representations refer to holistic or global structures, not separate properties of an object which can be listed individually or as a set of propositions.

Criticisms concerning the use of analog representations as explanatory constructs indicated the incoherence of the "picture metaphor" for mental imagery (Shepard & Cooper, 1982). Also, these emphasized the necessity for proposing a single basic symbolic representa-

tional system for translating between spatial and verbal processes (Pylyshyn, 1977). Other criticisms (Anderson, 1978) claimed that propositional processes and representations can mimick phenomena normally employed to support the requirement for analog representations and processes. Mental transformations and rotations have been considered to be primarily analog processes (e.g., Attneave, 1974; Cooper & Shepard, 1973; Kosslyn, 1980; Kosslyn & Pomerantz, 1977; Paivio, 1976). This assertion is considerably contentious since it seems to undermine the notion that human cognition can be simulated by a digital computer (Corballis, 1982). The study of mental rotation and other kinds of spatial transformation may necessitate a change in the models that are typically constructed, e.g., network theories of cognition (Anderson & Bower, 1973; Norman, Rumelhart & LNR Research Group, 1975).

A number of experiments have demonstrated that mental rotation of spatial objects seems to occur in real time. Visualizing or imagining a shape in a different orientation is tantamount to actually seeing it in that position. Apparently, mental rotation is similar to, or interacts with, perceived rotation. The findings of some investigations have confirmed the belief that mental rotation is a global, holistic, uniform, and continuous process which approximates to some extent actual physical rotation. That is, rotating a shape cognitively is analogous to rotating it realistically. This seems to imply a similarity in neurophysiological processes which take place when one is imagining or perceiving a spatial transformation or rotation (Corballis, 1982; Shepard & Chipman, 1970; Shepard & Cooper, 1982).

Introspective reports from subjects have indicated that they did imagine or visualize when performing mental transformations in the form of a "paper folding" or "surface development" task. These paper-and-pencil tests (Ekstrom, French, Harman, & Derman, 1976; Smith, McFarlane, 1964) of mental transformations have been used for a considerable length of time for assessing spatial ability. Paper folding or surface development tasks can be considered as mental rotation tasks. That is, each mental fold can be regarded as a rotation. However, each cognitive fold is conceived of as a rigid rotation of only one segment of the object relative to the rest--not the rotation of the complete object. Also, an individual mental rotation or fold

can be thought of as only one in a sequence that must be completed before a subject can give a response (Shepard & Cooper, 1982).

Measures of cerebral electrical activity, electroencephalographic (EEG) and event-related potential (ERP) recordings, represent this neurophysiological process as minute signals obtained from the scalp. The EEG depicts on-going brain activity while the ERP depicts this activity after specific stimulus events, e.g., light flashes or audible clicks. Typically, for people performing propositional or verbal tasks such as reading prose passages, there is decreased activity over the left hemisphere. For analog or spatial tasks such as recognizing random shapes, there is generally a decrease in activity over the right hemisphere. Such decreases may be considered indices of increased information processing within the affected hemisphere. That is, at least two modes of cognitive processing have been shown to be related to the brain's two hemispheres. A propositional, verbal, analytic, sequential, logical manner of information processing has been associated with left- hemisphere activity in most right-handed individuals. Conversely, an analog, spatial, holistic, simultaneous, intuitive manner has been attributed to right hemisphere activity (e.g., Allen, 1983; Bryden, 1982; Helligie, 1980; Hilyard & Kutas, 1983). There is considerable consensus concerning the existence of cerebral lateralities; however, their specific contributions to particular cognitive functions are still very much debatable and continue to be the object of scientific investigation.

It seems reasonable for auditory event-related potentials to trigger, elicit, evoke, or engage the neurophysiological correlates of propositional processing and representation. Likewise, it seems reasonable for visual event-related potentials to trigger, elicit, evoke, or engage the neurophysiological correlates of analog processing and representation. When auditory and visual stimuli are presented together, bimodal event-related potentials, then both propositional and analog processes and representations should be triggered or elicited. This assumes that perceptual processes and neurophysiological correlates of representational systems are approximately equivalent or at least interact.

Sensory interaction (Butterworth, 1981; Lewis & Froning, 1981; Shipley, 1970, 1977; Ship-

ley & Jones, 1969; Walk & Pick, 1981) has been considered as a joint function of auditory, visual, and bimodal event-related potentials, i.e., AERP, VERP, and BERP, respectively. As a neurophysiological index it has been conceptualized in terms of the absolute values of BERP and (AERP + VERP). If BERP is larger than (AERP + VERP), this has been thought of as an indicator of neurophysiological excitation or facilitation in cerebral activity. If BERP is smaller than (AERP + VERP), this has been thought of as an indicator of neurophysiological inhibition or suppression in cerebral activity. Since the neurophysiological correlates of sensory, perceptual, and representational processes are assumed to be tantamount or at least intermesh, sensory interaction then reflects the interrelationship, interplay, or interassociation of analog and propositional representational systems. Consequently, with BERP larger than (AERP + VERP) these dual representational processes support, reinforce, or correspond to one another. With BERP smaller than (AERP + VERP) they undermine, weaken, or nonconform to one another.

Imaginal and verbal encoding systems or representational modes have been hypothetically implicated as mediation mechanisms employed in the performance of spatial tasks (Paivio, 1971). The imaginal, visual, or analog encoding or representational system is primarily a parallel processing one specialized for storage and symbolic manipulation of spatial objects; whereas, the verbal, auditory, or propositional encoding or representational system is primarily a sequential processing one because of its linguistic nature. These representational mechanisms do not normally function independently of one another or only in one capacity. They are thought to interact continuously in performing complicated spatial tasks.

Imaginal and verbal encoding mechanisms can either facilitate or inhibit the performance of spatial ability tasks. Facilitation should occur where both representational systems are congruent with the processes demanded by the task; inhibition should occur where either or both of these symbolic systems are not congruent with those processes. Visual and auditory sensory input does impact upon intricate information processing. The sensory interaction or integration between these two stimulus modalities should become increasingly important as

higher-order cognitive functioning is required by the task. If sensory inputs, perceptual processes, and representational operations involve some of the same cerebral mechanisms, then how the brain integrates visual and auditory stimuli resulting in facilitation or inhibition should be related to its performance on complex spatial tasks. This is so since the visual-imaginal-analog system and the auditory-verbal- propositional system are speculatively implicated in spatial ability.

The objectives of this research were to (a) ascertain the amount of association between cerebral sensory interaction and spatial ability, (b) determine the importance of the visual-imaginal-analog system as well as the verbal- auditory-propositional system for spatial task performance, and (c) provide converging support that the proper interaction or integration of these dual representational systems is associated with spatial ability.

Method

Subjects

The subjects were fifty, right-handed, male, Caucasian recruits from the Naval Training Center, San Diego, who were undergoing basic enlisted military instruction. Audition and vision of the subjects were empirically verified as normal.

Spatial Ability Measure

The Surface Development Test (Ekstrom, French, Harman, & Derman, Note 1) was employed as a measure of spatial ability. This is a paper-and-pencil test of mental "paper folding" and "surface development". Subjects are required to imagine or visualize how a piece of paper can be folded to form some object. Two drawings are presented for each item. The drawing on the left is of a piece of paper which can be folded on the dotted lines to form the object depicted on the right. The task demands that the subjects imagine the folding and figure out which of the lettered edges on the object are the same as the numbered edges on the piece of paper at the left. Figure 1 depicts a sample item from this test for the reader's inspection.

Insert Figure 1 about here

Subjects' scores on this test are the number of correct letters minus a fraction of the number of incorrect letters. They had six minutes to complete each of the two parts of this test which was administered to each subject counterbalanced with brainwave recording.

Instrumentation

Data were acquired on a field-portable computer system that included a Data General NOVA 2/10 central processing unit (CPU, 32K memory); a dual-drive floppy disk unit (Advanced Electronics Design, Inc., Model 2500); an optically isolated and multiplexing EEG unit with band pass set for 0.2-30 Hz; signal sampling rate for each channel of the analogue EEG waveforms was set at 500 Hz; and a videographic display unit, integrated into the CPU, that presented visual stimuli to the subjects and displayed the analyzed ERP data. Permanent storage of all video information was obtained from a video hard copy unit (Tektronix Model 4632).

Stimuli

Visual (V) stimuli were computer-generated black and white checker-board patterns presented over the video monitor (Panasonic 14-inch Model WV 5400). Binocular visual field stimulation was about 9 degrees visual angle. Each check subtended about 17 minutes visual angle. Average background luminance was about 0.3 ftL and target luminance was about 5 ftL. The patterns were presented aperiodically, with interstimulus intervals averaging about 2 seconds (1.0-3.0 seconds).

Auditory (A) clicks were presented binaurally over headphones (Sennheier Model 424X) aperiodically about every 2 seconds (1.0-3.0 second interstimulus intervals. Click intensity was about 65 dB (A) (Bruel and Kjaer Impulse Sound Level Meter, Model 2209, One-Third Octave Filter Set, Model 1616). Headphone leads were shielded to minimize click artifacts.

Bimodal (B) presentation included simultaneous presentation of the visual and auditory stimuli. These stimuli were presented aperiodically about every 2 seconds (1.0-3.0 seconds).

During all recording periods, white noise was used for masking. It was presented to the subjects through the headphones and via a speaker in the sound chamber at a level of approximately 50 db (A). This was done to create more uniform data acquisition conditions across all the subjects. The auditory click stimuli were presented over this background noise.

Procedure

Recording Sites

Eight channels of visual, auditory, and bimodal ERP data were acquired from four pairs of homologous sites. Sites F3 and F4 are over the frontal brain region, an association area; sites T3 and T4 are over the temporal region, a primary auditory reception area where many visual and auditory nerves interconnect; sites P3 and P4 are over the parietal region, a primary association area; and sites 01 and 02 are over the occipital region, a primary visual reception area. Ground was at Pz in the mid-parietal area. Sites designated by odd numbers denote left hemisphere locations; and those designated by even numbers, right hemisphere locations.

Electrodes

The subjects were prepared for recording after they had received brief instruction. They completed a brief background questionnaire and signed a privacy act and volunteer consent form. An elastic helmet (Lycra) fitted with plastic holders for the electrodes was placed on the subject's head. Each subject's hair was parted and scalp cleaned with an alcohol-impregnated cotton swab that was placed through the holders. Electrode cream was placed down the holders and rubbed into the scalp. The electrodes were silver/silver chloride Beckman miniatures (actual surface contact area was 2mm) with a clear plastic extension tube (38mm long) attached and filled with electrolytic solution. A small sponge (microcell foam) soaked with electrolyte held the solution in the tube and made contact with the electrode paste on the scalp. The extension tube not only held the electrode in place but also minimized

the slow potential drift due to scalp temperature change that would have otherwise been picked up at the recording site. A Beckman mini-electrode fitted with a standard two-sided adhesive wafer served as a reference electrode on the nose.

The helmet and all 10 electrodes could be attached in 6-8 minutes with impedance readings of 2-3K ohms. After all electrodes were in place, the subjects were instructed to observe their real-time EEG activity on the oscilloscope display. They were then instructed to move their jaws, eyebrows, etc., so that they could observe how muscle artifacts could contaminate the ERP data. The subject was then seated in a sound chamber in alignment with the video monitor. A hand-held switch allowed the subject to suspend all stimulus presentation and analysis operations to eliminate artifact. Additional artifact rejection was available by the console operater prior to storing the data.

Event-Related Potential (ERP) Data

ERPs were generated on-line, and the A/D digitization rate was one-quarter million samples per second. The visual and auditory ERP data were retrieved from a floppy diskette and the required computations were performed. The data were then displayed on the video monitor and hard copies were obtained. Bimodal ERP data were also computed and displayed in a similar manner.

Eight channels of visual and auditory ERP data were overlaid in Figure 2. Root mean square (RMS) and standard deviation (SD) amplitude values are presented, along with the waveform means values for the half-second post-stimulus epoch (533 msec). SD amplitude values (in microvolts (uV)) are normalized (waveform mean set to zero) RMS values (in uV). For all analyses, only SD amplitude values (in uV) were used. These values have been found to be very effective when employing ERPs to study individual differences among many different kinds of subjects. There are individuals who do not manifest clearly defined ERP components. This index is also advantageous in that it permits the description of ERP amplitude as a single value (Callaway, 1975; Lewis & Froning, 1981). Prestimulus waveforms (133 msec) were also recorded and displayed for each channel. Calibration, polarity, DC offset,

time base, and other descriptive information were also displayed. The waveforms in the left column were derived from the left hemisphere (LH). The waveforms from top to bottom were from the front to the back of the head at frontal, temporal, parietal, and occipital sites (F3, T3, P3, O1). Right hemisphere (RH) ERP data from sites F4, T4, P4, and O2 were represented in the right column.

Insert Figure 2 about here

Sensory Interaction Indices

In addition to the directly recorded ERP amplitudes, an index was derived to reflect an hypothesized property of brain behavior. In order to appraise sensory interaction and its association with spatial ability, the relationship between single modality presentations (visual or auditory) and the joint bimodal presentation (simultaneous stimulation of visual checkerboard pattens and auditory clicks) was determined. A sensory interaction index was defined as [BERP - (VERP + AERP)] (Federico, Froning, Calder, Note 2; Lewis & Froning, 1981; Lewis, Federico, Froning, & Calder, Note 3; Shipley, 1970; Shipley & Jones, 1969). If the result of this expression is positive, then the bimodal stimulation presentation produces a greater magnitude response than the sum of separate visual or auditory stimulation. This indicates sensory excitation, facilitation, or potentiation in the brain's nervous system. If the expression is negative, the sensory attenuation, inhibition, or suppression occurs in the nervous system when auditory stimulation is presented at the same time as visual stimulation. This sensory interaction index was derived for left and right hemisphere frontal, temporal, parietal, and occipital areas.

Statistical Analysis

Product-moment correlations were computed between sensory interaction indices for left and right hemisphere sites and Surface Development Test scores. Means and standard devia-

tions were also computed for these measures.

Results

The means and standard deviations for left and right hemisphere sensory interactions, together with the correlations between these derived indices and spatial ability test scores ($\bar{X}$ = 28.34; SD = 16.76), are presented in Table 1. The negative means for the sensory interaction indices indicated that

---

Insert Table 1 about here

---

attenuation, inhibition, or suppression occurred at all cerebral areas. The standard deviations revealed that this attenuated activity appeared approximately uniform between homologous hemispheric sites. The product-moment correlations disclosed that sensory interaction was positively related to spatial ability. Sensory interaction indices for all four left hemisphere sites (F3, T3, P3, and O1) were significantly associated with scores on the Surface Development Test; and sensory interaction indices for three right hemisphere sites (F4, T4, and P4) were significantly associated with scores on the Surface Development Test. The above statistics implied that sensory suppression, inhibition, or attenuation of the cortex is significantly related to spatial ability. The nature of this relationship is as follows: as sensory interaction indices become less negative, spatial ability scores become more positive. That is, as sensory suppression lessens, spatial ability increases.

Discussion and Conclusions

This study was scientifically salient since: (a) it reports correlations between cerebral sensory interaction and spatial ability, (b) it implicates the visual-imaginal-analog system as well as the auditory-verbal-propositional system in spatial task performance, and (c) it provides converging support that the proper interaction or integration between these dual representa-

tional systems is associated with spatial ability.

The results indicated that cerebral sensory interaction is positively related to spatial ability. How the brain integrates visual and auditory stimuli is associated with performance on a complex spatial task. Sensory inhibition, attenuation, or suppression at left and right frontal, temporal, and parietal areas was positively correlated with spatial ability. Not only the spatially superior and parallel processing right hemisphere but also the linguistically superior and linear processing left hemisphere is associated with spatial task performance. The findings supported the theory that the visual-imaginal- analog system and the auditory-verbal-propositional system are involved in spatial ability. The extent to which the cortex can suppress, inhibit, or attenuate the interaction or integration between these dual representational systems is related to spatial task performance.

In regards to the dispute dealing with the nature of the mental representations inherent to spatial ability, the results suggested that another particular genre of cognition, analog processing, is required in addition to that associated with propositional processing. It was established that as sensory suppression lessens, spatial ability increases. This seemed to imply that visual and auditory input impacts upon intricate information processing, and that both representational systems undoubtedly influence complex spatial performance. As higher-order cognitive functioning is demanded by the spatial task, sensory interaction between these two stimulus modalities appears to become increasingly important. How the brain integrates visual and auditory stimuli culminating in sensory suppression seems to be associated with performance on complicated spatial tasks. It was assumed that the neurophysiological correlates of sensory, perceptual, and representational processes are equivalent or at least intermesh. Therefore, sensory interaction reflects the interrelationship, interplay, or interassociation of analog and propositional representational systems. Consequently, the results speculatively implicate both of these representational processes in spatial ability.

In agreement with Anderson (1978), who has argued that analog and propositional hypotheses cannot be differentiated on the basis of behavioral evidence, this study insinuated

that these dual symbol systems are intimated in spatial reasoning. It also involved the left and right cerebral hemisphere in this ability. During this research, paper folding or surface development was considered to be a mental rotation task. Each mental fold was regarded as a rigid rotation of one part of the object relative to the rest. Assuming that mental rotation is a primary component of spatial processing, it would typically and chiefly be expected to be a right hemisphere task. This is especially so in view of this hemisphere's well documented superiority in the execution of spatial skills (e.g., Allen, 1983; Bryden, 1982; Helligie, 1980; Hillyard & Kutas, 1983) Most, but not all, the evidence corroborate this expectancy (e.g., Benton, 1983; Newcombe, 1983; Ratcliff, 1983).

Contrary to the norm, a study by Ornstein, Johnstone, Herron, and Swencionis (1980) established that more electroencephalographic activity occurred over the "left" than the right cerebral hemisphere when subjects performed a mental rotation task. De Renzi (1978) reviewed other evidence that implicated the "left" hemisphere in the execution of complex spatial tasks. He summarized the findings by concluding that the more complicated the spatial task, the more involved the "left" hemisphere. This is consonant with data obtained by others (e.g., Federico, 1980a, 1980b; Federico & Montague, 1975) that suggested difficult spatial performance was facilitated by verbal, semantic, or propositional representations or encodings. This kind of mental mediation has been primarily linked with the left hemisphere (e.g., Allen, 1983; Bryden, 1982; Helligie, 1980; Hillyard & Kutas, 1983). As implied by this reported research, surface development being a complex spatial task taxes the respective resources of the left and right cerebral hemispheres: analog as well as propositional representational systems. To associate spatial ability with either hemisphere exclusively or chiefly would be misleading. Caution must also be exercised against absolutely attributing analog symbolizing processes to the right hemisphere, and propositional processes to the left hemisphere.

This does not preclude the consideration of the visual, imaginal, or analog symbolizing system as a distinct or separate representational structure which uses a "quasi-pictorial format" (Kosslyn, 1980). Individual differences in analog representational processes and structures, as

well as complementary propositional systems, can contribute to individual differences in spatial ability (Lohman & Kyllonen, 1983). One hypothesis proposes that it is probably the capacity of the visual short-term memory or buffer, which determines the accuracy and rapidity to generate and refresh images, that accounts for a considerable amount of the variability in spatial ability (Kosslyn, 1981). It is possible that subjects who were unable, or found it difficult, to conjure up a mental image of each object while performing the surface development task were at a certain disadvantage. If these subjects were engaging exclusively or chiefly their verbal, semantic, or propositional representational system when attempting the paper folding task, then they were imposing an incompatible cognitive style of processing which probably prevented successful spatial performance by shunting their visual short-term-memory system. It seems likely that subjects who performed the surface development task well, thus scoring high in spatial ability, had sufficient sensory suppression or inhibition to attenuate the verbal or propositional representational system. This would enable them to more completely and properly employ the visual short-term-memory buffer, and its corresponding imaginal or analog representational system, to optimize the performance of the spatial task. This is not meant to imply, however, that the propositional representational system was entirely "out of the picture".

Assuming that spatial proficiency involves sensual, perceptual, and representational processes and that the neurophysiological correlates of these cortical operations are tantamount or at least intermesh, this study examined this cognitive ability by associating subjects' performance on a marker test for this intellectual skill with cerebral indices of their sensory interaction. Other researchers have used sensory interaction indices to investigate different mental phenomenon. Shipley (1970), who employed the sensory interaction approach to study retarded children, reported that they were deficient chiefly in their capacity to attend to and monitor multidimensional stimuli. Using similar methodology, Shipley and Jones (1969) that found dyslexics could not process bimodally presented sensory input as well as normal children. They asserted that normals manifested more frequently sensory facilitation or excita-

tion than dyslexics. Also, these latter children demonstrated some modest amounts of attenuation, or at least, lack of sensory arousal. Barnet and Lodge (1967), however, reported probable defects of sensory inhibitory mechanisms in mongoloids culminating in more pronounced response habituation in mongoloids than in normals. They speculated that such a loss of suppressing mechanisms in retardates could account for their extreme perseveration in excessively monotonous tasks. Lewis and Froning (1981) found that subjects with high reading ability demonstrated more sensory inhibition than those with low. They maintained that such suppression or attenuation is usually associated with normal cortical processes. The major result of this research was: as sensory suppression lessens, spatial ability increases. Consequently, this study's salient finding is completely consonant with the results of Barnet and Lodge (1967), Lewis and Froning (1981), and Shipley (1970), and only partially compatible with the outcome of Shipley and Jones (1969).

Reference Notes

[1] Ekstrom, R. B., French, J. W., Harman, H. H., & Derman, D. *Manual for kit of factor-referenced cognitive tests*. Princeton, NJ: Educational Testing Service, 1976.

[2] Federico, P-A., Froning, J. N., & Calder, M. *Validation of brain event-related potentials as indicators of cognitive styles, abilities, and aptitudes (NPRDC TR 83-11)*. San Diego, CA: Navy Personnel Research and Development Center, 1983.

[3] Lewis, G. W., Federico, P-A., Froning, J. N., & Calder, M. *Event-related brain potentials and cognitive processing: Implication for navy training (NPRDC TR 82-8)*. San Diego, CA: Navy Personnel Research and Development Center, 1981.

References

Allen, M. Models of hemispheric specialization. *Psychological Bulletin*, 1983, *93*, 73-104.

Anderson, J. R. Arguments concerning representations for mental imagery. *Psychological Review*, 1978, *85*, 249-277.

Anderson, J. R., & Bower, G. H. *Human associative memory*. Washington, DC: Hemispheric Press, 1973.

Attneave, F. How do you know? *American Psychologist*, 1974, *29*, 493-499.

Barnet, A. B., & Lodge, A. Click evoked EEG responses in normal and developmentally retarded infants. *Nature*, 1967, *214*, 252-255.

Benton, A. Spatial thinking in neurological patients: Historical aspects. In M. Potegal (Ed.), *Spatial abilities: Development and physiological foundations*. New York: Academic Press, 1983.

Bryden, M. P. *Laterality: Functional asymmetry in the intact brain*. New York: Academic Press, 1982.

Butterworth, G. The origins of auditory-visual perception and visual proprioception in human development. In R. D. Wick & H. L. Pick, Jr. (Eds.), *Intersensory perception and sensory integration*. New York: Plenum Press, 1981.

Cooper, L. A., & Shepard, R. N. Chronometric studies of the rotation of mental images. In W. G. Chase (Ed.), *Visual information processing*. New York: Academic Press, 1973.

Corballis, M. C. Mental rotation: Anatomy of a paradigm. In M. Potegal (Ed.), *Spatial ability: Development and physiological foundations*. New York: Academic Press, 1982.

De Renzi, E. Hemispheric asymmetry as evidenced by spatial disorders. In M. Kinsbourne (Ed.), *Asymmetrical function of the human brain*. New York: Academic Press, 1978.

Federico, P-A. Some effects of encoding, codability, and exposure upon recognition memory. *Bulletin of the Psychonomic Society*, 1980, *16*, 89-92. (a)

Federico, P-A. Domain of processing and recognition memory for shapes. *Bulletin of the Psychonomic Society*, 1980, *16*, 261-264. (b)

Federico, P-A., & Montague, W. E. Recognition memory as a function of encoding strategy and stimulus codability. *Journal of Experimental Psychology: Human Learning and Memory*, 1975, *1*, 680-688.

Helligie, H. H. Cerebral hemisphere asymmetry: Methods, issues, and implications. *Educational Communication and Technology Journal*, 1980, *28*, 83-98.

Hermelin, B., & O'Conner, N. Spatial modality coding in children with and without impairments. In M. Potegal (Ed.), *Spatial abilities: Development and physiological foundations*. New York: Academic Press, 1982.

Hillyard, S. A., & Kutas, M. Electrophysiology of cognitive processing. *Annual Review of Psychology*, 1983, *34*, 33-61.

Kosslyn, S. M. *Image and mind*. Cambridge, MA: Harvard University Press, 1980.

Kosslyn, S. M. The medium and the message in mental imagery: A theory. *Psychological Review*. 1981, *88*, 46-66.

Kosslyn, S. M. & Pomerantz, J. R. Imagery, propositions, and the form of internal representations. *Cognitive Psychology*, 1977, *9*, 52-76.

Lewis, G. W., & Froning, J. N. Sensory interaction, brain activity, and reading ability in young adults. *International Journal of Neuroscience*, 1981, *15*, 128-140.

Lohman, D. F., & Kyllonen, P. C. Individual differences in solution strategy on spatial tasks. In R. F. Dillon & R. R. Schmeck (Eds.), *Individual differences in cognition, Volume 1*. New York: Academic Press, 1983.

Newcombe, N. Sex-related differences in spatial ability: Problems and gaps in current approaches. In M. Potegal (Ed.), *Spatial abilities: Development and physiological foundations*. New York: Academic Press, 1982.

Norman, D. A., & Rumelhart, D. E., & the LNR Research Group. *Explorations in cognition*. San Francisco: Freeman, 1975.

Ornstein, R., Johnstone, J., Herron, J., & Swencionis, C. Differential right hemisphere engagement in visuospatial tasks. *Neuropsychologia*, 1980, *18*, 49-64.

Paivio, A. *Imagery and verbal processes*. New York: Holt, Rinehart and Winston, 1971.

Paivio, A. Images, propositions, and knowledge. In J. M. Nicholas (Ed.), *Images, perception, and knowledge*. Dorecht, The Netherlands: Reidel, 1976.

Pylyshyn, Z. W. The rate of "mental rotation" of images: A test of a holistic analogue hypothesis. *Memory and Cognition*, 1979, *7*, 19-28.

Ratcliff, G. Disturbances of spatial orientation associated with cerebral lesions. In M. Potegal (Ed.), *Spatial abilities: Development and physiological foundations*. New York: Academic Press, 1982.

Shepard, R. N., & Chipman, S. Second-order isomorphism of internal representations: Shapes of states. *Cognitive Psychology*, 1970, *1*, 1-17.

Shepard, R. N., & Cooper, L. A. *Mental images and their transformation*. Cambridge, MA: The MIT Press, 1982.

Shipley, T. Evoked brain potentials and sensory interaction in the retarded child. *American Journal of Mental Deficiency*, 1970, *74*, 517-523.

Shipley, T., & Jones, R. W. Initial observations on sensory interaction and the theory of

dyslexia. *Journal of Communication Disorders*, 1969, 2, 295-311.

Shipley, T. Interhemispherical differences in the evoked cortical potential to intersensory and repetitive stimulation: Hypotheses, methods, and appraisals. *Neuropsychologia*, 1977, *15*, 133-141.

Smith, I. McFarlane. *Spatial ability*. London: University of London Press, 1964.

Walk, R. D. & Pick, Jr., H. L. *Intersensory perception and sensory integration*. New York: Plenum Press, 1981.

Footnotes

Opinions and assertions contained herein are those of the author and are not to be construed as official or reflecting the views of the Department of the Navy or Defense.

Requests for reprints should be sent to Pat-Anthony Federico, Code P51, NPRDC, San Diego CA 92152.

The author acknowledges the assistance of Gregory W. Lewis and Jeffrey N. Froning during the data collection phase of this research.

Table 1

Descriptive Statistics for Sensory Interactions and their Correlations with Spatial Ability

Sensory Interaction	X	SD	r
Left Hemisphere			
Frontal	-0.96	2.01	.43**
Temporal	-1.61	1.46	.31*
Parietal	-1.75	1.54	.36**
Occipital	-2.01	1.71	.37**
Right Hemisphere			
Frontal	-1.21	2.01	.34*
Temporal	-1.57	1.25	.37**
Parietal	-1.81	1.42	.28*
Occipital	-1.55	1.50	.26

*$r(48) \geq .28$; $p < .05$.
**$r(48) \geq .36$; $p < .01$.

Figure Captions

Figure 1. Sample item from the Surface Development Test used to measure spatial ability.

Figure 2. Sample visual and auditory event-related potential data. The left column is from the left hemisphere; and the right column, from the right hemisphere. From top to bottom, the records are from the frontal, temporal, parietal, and occipital areas.


2
4
1
X
5
3
E
H
B
F
C
A
X
D
G
1:
2:
3:
4:
5:


REF
(ANTERIOR)
[REFER TO NOSE]
[SCALE =100.00 Z]
** VERP TOPOGRAPHY **
[OVERLAY]
1ST 10 VISUAL
ALL 10 AUDITORY .....
[BOTH EARS]
POST-STIM (UV)
(UN-NORMED)
2.96 F3
1.61
2.49
F4
3.95
1.86
3.48
(+)
5 UV
(-)
RMS= 4.01 T3
SD = 2.43
MN = 3.19
T4
3.46
2.26
2.62
L
R
1.19 P3
1.17
0.23
P4
3.00
1.39
2.66
3.25 O1
3.20
0.52
O2
3.62
3.60
0.36
0.0
STIM
-133.33
533.33
MSEC
(POSTERIOR)
(POST-STIM MEANS = 0.0)